


# Applications of Forces - Questions

---

May 2013 Mathematics Advanced Paper 1: Mechanics 1

- 1.
- 3.




**Figure 1**

A box of mass 2 kg is held in equilibrium on a fixed rough inclined plane by a rope. The rope lies in a vertical plane containing a line of greatest slope of the inclined plane. The rope is inclined to the plane at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$ , and the plane is at an angle of  $30^\circ$  to the horizontal, as shown in Figure 1. The coefficient of friction between the box and the inclined plane is  $\frac{1}{3}$  and the box is on the point of slipping up the plane. By modelling the box as a particle and the rope as a light inextensible string, find the tension in the rope.

**(8)**

2.

3.



**Figure 2**

A box of mass 5 kg lies on a rough plane inclined at  $30^\circ$  to the horizontal. The box is held in equilibrium by a horizontal force of magnitude 20 N, as shown in Figure 2. The force acts in a vertical plane containing a line of greatest slope of the inclined plane.

The box is in equilibrium and on the point of moving down the plane. The box is modelled as a particle.

Find

(a) the magnitude of the normal reaction of the plane on the box,


(4)

(b) the coefficient of friction between the box and the plane.

(5)

3.

8.



**Figure 2**

A particle  $P$  of mass  $4\text{ kg}$  is moving up a fixed rough plane at a constant speed of  $16\text{ m s}^{-1}$  under the action of a force of magnitude  $36\text{ N}$ . The plane is inclined at  $30^\circ$  to the horizontal. The force acts in the vertical plane containing the line of greatest slope of the plane through  $P$ , and acts at  $30^\circ$  to the inclined plane, as shown in Figure 2. The coefficient of friction between  $P$  and the plane is  $\mu$ . Find

(a) the magnitude of the normal reaction between  $P$  and the plane, (4)

(b) the value of  $\mu$ . (5)

The force of magnitude  $36\text{ N}$  is removed.

(c) Find the distance that  $P$  travels between the instant when the force is removed and the instant when it comes to rest. (5)

4.  
3.




Figure 1

A particle of weight  $W$  newtons is held in equilibrium on a rough inclined plane by a horizontal force of magnitude 4 N. The force acts in a vertical plane containing a line of greatest slope of the inclined plane. The plane is inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$  as shown in Figure 1.

The coefficient of friction between the particle and the plane is  $\frac{1}{2}$ .

Given that the particle is on the point of sliding down the plane,

- (i) show that the magnitude of the normal reaction between the particle and the plane is 20 N,  
(ii) find the value of  $W$ .

(9)

5.  
6.




Figure 2

Two particles  $P$  and  $Q$  have masses 0.3 kg and  $m$  kg respectively. The particles are attached to the ends of a light inextensible string. The string passes over a small smooth pulley which is fixed at the top of a fixed rough plane. The plane is inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$ . The coefficient of friction between  $P$  and the plane is  $\frac{1}{2}$ .

The string lies in a vertical plane through a line of greatest slope of the inclined plane. The particle  $P$  is held at rest on the inclined plane and the particle  $Q$  hangs freely below the pulley with the string taut, as shown in Figure 2.

The system is released from rest and  $Q$  accelerates vertically downwards at  $1.4 \text{ m s}^{-2}$ .

Find

(a) the magnitude of the normal reaction of the inclined plane on  $P$ , (2)

(b) the value of  $m$ . (8)

When the particles have been moving for  $0.5 \text{ s}$ , the string breaks. Assuming that  $P$  does not reach the pulley,

(c) find the further time that elapses until  $P$  comes to instantaneous rest. (6)

Jan 2011 Mathematics Advanced Paper 1: Mechanics 1

6.

6.




Figure 4

A particle of weight  $120 \text{ N}$  is placed on a fixed rough plane which is inclined at an angle  $\alpha$  to the horizontal, where  $\tan \alpha = \frac{3}{4}$ .

The coefficient of friction between the particle and the plane is  $\frac{1}{2}$ .

The particle is held at rest in equilibrium by a horizontal force of magnitude  $30 \text{ N}$ , which acts in the vertical plane containing the line of greatest slope of the plane through the particle, as shown in Figure 2.

- (a) Show that the normal reaction between the particle and the plane has magnitude 114 N. (4)




Figure 3

The horizontal force is removed and replaced by a force of magnitude  $P$  newtons acting up the slope along the line of greatest slope of the plane through the particle, as shown in Figure 3. The particle remains in equilibrium.

- (b) Find the greatest possible value of  $P$ . (8)
- (c) Find the magnitude and direction of the frictional force acting on the particle when  $P = 30$ . (3)

7.  
7.




Figure 4

Two particles  $A$  and  $B$ , of mass  $7\text{ kg}$  and  $3\text{ kg}$  respectively, are attached to the ends of a light inextensible string. Initially  $B$  is held at rest on a rough fixed plane inclined at angle  $\theta$  to the horizontal, where  $\tan \theta = \frac{5}{12}$ . The part of the string from  $B$  to  $P$  is parallel to a line of greatest slope of the plane. The string passes over a small smooth pulley,  $P$ , fixed at the top of the plane. The particle  $A$  hangs freely below  $P$ , as shown in Figure 4. The coefficient of friction between  $B$  and the plane is  $\frac{2}{3}$ . The particles are released from rest with the string taut and  $B$  moves up the plane.

(a) Find the magnitude of the acceleration of  $B$  immediately after release. (10)

(b) Find the speed of  $B$  when it has moved  $1\text{ m}$  up the plane. (2)

When  $B$  has moved  $1\text{ m}$  up the plane the string breaks. Given that in the subsequent motion  $B$  does not reach  $P$ ,

(c) find the time between the instants when the string breaks and when  $B$  comes to instantaneous rest. (4)

May 2010 Mathematics Advanced Paper 1: Mechanics 1

8.

3.





Figure 1

A small box is pushed along a floor. The floor is modelled as a rough horizontal plane and the box is modelled as a particle. The coefficient of friction between the box and the floor is  $\frac{1}{2}$ . The box is pushed by a force of magnitude  $100\text{ N}$  which acts at an angle of  $30^\circ$  with the floor, as shown in Figure 1.

Given that the box moves with constant speed, find the mass of the box. (7)

9.  
7.



**Figure 2**

A particle of mass  $0.4$  kg is held at rest on a fixed rough plane by a horizontal force of magnitude  $P$  newtons. The force acts in the vertical plane containing the line of greatest slope of the inclined plane which passes through the particle. The plane is inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$ , as shown in Figure 2.

The coefficient of friction between the particle and the plane is  $\frac{1}{3}$ .

Given that the particle is on the point of sliding up the plane, find

(a) the magnitude of the normal reaction between the particle and the plane,

(5)

(b) the value of  $P$ .

(5)

10.

5. A particle of mass  $0.8 \text{ kg}$  is held at rest on a rough plane. The plane is inclined at  $30^\circ$  to the horizontal. The particle is released from rest and slides down a line of greatest slope of the plane. The particle moves  $2.7 \text{ m}$  during the first  $3 \text{ seconds}$  of its motion. Find

(a) the acceleration of the particle,

(3)

(b) the coefficient of friction between the particle and the plane.

(5)

The particle is now held on the same rough plane by a horizontal force of magnitude  $X \text{ newtons}$ , acting in a plane containing a line of greatest slope of the plane, as shown in Figure 3. The particle is in equilibrium and on the point of moving up the plane.




Figure 3

(c) Find the value of  $X$ .

(7)

11.

6.




Figure 4

Two particles  $A$  and  $B$  have masses  $5m$  and  $km$  respectively, where  $k < 5$ . The particles are connected by a light inextensible string which passes over a smooth light fixed pulley. The system is held at rest with the string taut, the hanging parts of the string vertical and with  $A$  and  $B$  at the same height above a horizontal plane, as shown in Figure 4. The system is released from rest. After release,  $A$  descends with acceleration  $\frac{1}{4}g$ .

(a) Show that the tension in the string as  $A$  descends is  $\frac{15}{4}mg$ . (3)

(b) Find the value of  $k$ . (3)

(c) State how you have used the information that the pulley is smooth. (1)

After descending for 1.2 s, the particle  $A$  reaches the plane. It is immediately brought to rest by the impact with the plane. The initial distance between  $B$  and the pulley is such that, in the subsequent motion,  $B$  does not reach the pulley.

(d) Find the greatest height reached by  $B$  above the plane. (7)